Bivariate-Schurer-Stancu operators based on (p,q)-integers

Abdul Wafi, Nadeem Rao*

[a]*Department of Mathematics, Jamia Millia Islamia, New Delhi-110 025, India*

arXiv:1602.06315v1 [math.CA] 14 Jan 2016

Abstract

The aim of this article is to introduce a bivariate extension of Shurer-Stancu operators based on (p,q)-integers. We prove uniform approximation by means of Bohman-Korovkin type theorem, rate of convergence using total modulus of smoothness and degree of approximation by using second order modulus of smoothness, Peetre's K-functional, Lipschitz type class.

Keywords: Schurer-Stancu, (p,q)-integers, modulus of smoothness, rate of convergence.
2010 Mathematics Subject Classification 41A10, 41A25, 41A35, 41A36

1. Introduction

In 1962, Schurer [1] introduced the following generalization of the classical Bernstein operators for all non-negative integer l and $n \in \mathbb{N}$

$$B_n^l(f;x) = \sum_{k=0}^{n+l} \binom{n+l}{k} x^k (1-x)^{n+l-k} f\Big(\frac{k}{n}\Big),$$

where $f \in C[0, l+1]$ and $x \in [0,1]$.

In recent past, the applications of q-calculus attracted the attention of mathematicians and has an interesting impact in the research in approximation theory. It has been noticed that linear positive operators based on q-integers are quite effective as far as the rate of convergence is concerned. In 1987, Lupas [2] first defined q-analogue of Bernstein operators. In 1997, Philips [3] studied other form of Bernstein-polynomials based on q-integers. Several extensions of q-linear positive operators have been studied by different researchers for instance (see [4]). Recently, Mursaleen et al [5] added an idea based on (p,q)-calculus in approximation theory and gave a (p,q) extension to the classical Bernstein operators. The motive of (p,q)-integers was to generalize various forms of q-oscillator algebras in physics [6]. Several generalization of Bernstein operators were studied using (p,q)-analogue and their approximation properties have been investigated. For instance, (p,q)-Bernstein-Stancu operators [7]; (p,q)-Bernstein-Schurer operators [8]; (p,q)-Schurer-Stancu operators [9]; Chlodowski variant of (p,q)-Schurer-Stancu operators [10]; (p,q)-Szasz operators [11]; (p,q)-Szasz-Baskakov operators [12] were introduced and their approximation properties are studied. Motivated by the above generalizations, we present a bivariate extension of (p,q)-Schurer-Stancu operators in this paper.

Let $0 < q < p \leq 1$. Then, (p,q)-integers for non negative integers n,k are given by

$$[k]_{p,q} = \frac{p^k - q^k}{p-q} \qquad and \qquad [k]_{p,q} = 1 \qquad for \quad k = 0.$$

(p,q)-binomial coefficient

$$\binom{n}{k}_{p,q} = \frac{[n]_{p,q}!}{[k]_{p,q}![n-k]_{p,q}!},$$

and (p,q)-binomial expansion

$$\begin{aligned}(ax+by)_{p,q}^n &= \sum_{k=0}^{n} \binom{n}{k}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} a^{n-k} b^k x^{n-k} y^k,\\ (x+y)_{p,q}^n &= (x+y)(px+py)(p^2x+q^2y)...(p^{n-1}x-q^{n-1}y).\end{aligned}$$

*Corresponding author
Email addresses: awafi@jmi.ac.in (Abdul Wafi), nadeemrao1990@gmail.com (Nadeem Rao)

2. Construction of (p,q)-Bivariate-Schurer-Stancu operators

Let $I=[0,l+1]$ and $(x_1,x_2)\in I\times I=[0,l+1]\times[0,l+1]$. Then, for any function $f\in C(I\times I)$ and $(n_1,n_2)\in\mathbb{N}\times\mathbb{N}$, the operators $S_{n_{12},l}^{\alpha_{12},\beta_{12}}:C(I\times I)\to C([0,1]\times[0,1])$ is defined as follows

$$S_{n_{12},l}^{\alpha_{12},\beta_{12}}(f;p_{12},q_{12};x_1,x_2)=\sum_{\nu_1=0}^{n_1+l}\sum_{\nu_2=0}^{n_2+l}s_{n_1,l,\nu_1}^{p_1,q_1}(x_1)s_{n_2,l,\nu_2}^{p_2,q_2}(x_2)f\left(\frac{p^{n_1-\nu_1}[\nu_1]_{p_1q_1}+\alpha_1}{[n_1]_{p_1q_1}+\beta_1},\frac{p^{n_2-\nu_2}[\nu_2]_{p_2q_2}+\alpha_2}{[n_2]_{p_2q_2}+\beta_2}\right),\quad(2.1)$$

where $S_{n_{12},l}^{\alpha_{12},\beta_{12}}(f;p_{12},q_{12},;x_1,x_2)=S_{n_1,n_2,l_1,l_2}^{\alpha_1,\alpha_2,\beta_1,\beta_2}(f;p_1,q_1,p_2,q_2;x_1,x_2)$ and

$$s_{n_i,l,\nu_i}^{p_i,q_i}(x_i)=\frac{1}{p^{\frac{(n_i+l)(n_i+l-1)}{2}}}\binom{n_i+l}{\nu_i}_{p_i,q_i}p^{\frac{\nu_i(\nu_i-1)}{2}}x^{\nu_i}\prod_{j=0}^{n_i+l-\nu_i-1}(p_i^j-q_i^jx_i),$$

with the conditions
(i) for any positive real number p_i and q_i $(i=1,2)$ such that $0<q_i<p_i\le 1$,
(ii) for any non-negative real value of α_i and β_i $(i=1,2)$ such that $0\le\alpha_i\le\beta_i$.

Remark 2.1. One can find that
(i) if $p_i=1(i=1,2)$, then the operators defined by 2.1 reduce to q-Bivariate-Schurer-Stancu operators,

$$S_{n_{12},l}^{\alpha_{12},\beta_{12}}(f;q_1,q_2;x_1,x_2)=\sum_{\nu_1=0}^{n_1+l}\sum_{\nu_2=0}^{n_2+l}s_{n_1,l,\nu_1}^{q_1}(x_1)s_{n_2,l,\nu_2}^{q_2}(x_2)f\left(\frac{[\nu_1]_{q_1}+\alpha_1}{[n_1]_{q_1}+\beta_1},\frac{[\nu_2]_{q_2}+\alpha_2}{[n_2]_{q_2}+\beta_2}\right),$$

where

$$s_{n_i,l,\nu_i}^{q_i}(x_i)=\binom{n_i+l}{\nu_i}_{q_i}x^{\nu_i}\prod_{j=0}^{n_i+l-\nu_i-1}(1-q_i^jx_i),$$

(ii) if $\alpha_i=\beta_i=0,(i=1,2)$, then the operators defined by 2.1 reduce to (p,q)-Bivariate-Bernstein-Schurer operators

$$S_{n_{12},l}^{\alpha_{12},\beta_{12}}(f;p_{12},q_{12};x_1,x_2)=\sum_{\nu_1=0}^{n_1+l}\sum_{\nu_2=0}^{n_2+l}s_{n_1,l,\nu_1}^{p_1,q_1}(x_1)s_{n_2,l,\nu_2}^{p_2,q_2}(x_2)f\left(\frac{p^{n_1-\nu_1}[\nu_1]_{p_1q_1}}{[n_1]_{p_1q_1}},\frac{p^{n_2-\nu_2}[\nu_2]_{p_2q_2}}{[n_2]_{p_2q_2}}\right),$$

and
(iii) if $l_i=0$ and $\alpha_i=\beta_i=0,(i=1,2)$, then the operators defined by 2.1 reduce to (p,q)-Bivariate-Bernstein operators

$$S_{n_{12}}^{\alpha_{12},\beta_{12}}(f;p_{12},q_{12};x_1,x_2)=\sum_{\nu_1=0}^{n_1}\sum_{\nu_2=0}^{n_2}s_{n_1,\nu_1}^{p_1,q_1}(x_1)s_{n_2,\nu_2}^{p_2,q_2}(x_2)f\left(\frac{p^{n_1-\nu_1}[\nu_1]_{p_1q_1}}{[n_1]_{p_1q_1}},\frac{p^{n_2-\nu_2}[\nu_2]_{p_2q_2}}{[n_2]_{p_2q_2}}\right),$$

where

$$s_{n_i,\nu_i}^{p_i,q_i}(x_i)=\frac{1}{p^{\frac{(n_i)(n_i-1)}{2}}}\binom{n_i}{\nu_i}_{p_i,q_i}p^{\frac{\nu_i(\nu_i-1)}{2}}x^{\nu_i}\prod_{j=0}^{n_i-\nu_i-1}(p_i^j-q_i^jx_i).$$

Lemma 2.1. *Let $e_{i,j}=x_1^ix_2^j,0\le i,j\le 2$ are the two dimensional test functions. Then, we have*

$$\begin{aligned}
S_{n_{12},l}^{\alpha_{12},\beta_{12}}(e_{0,0};p_{12},q_{12};x_1,x_2) &= 1,\\
S_{n_{12},l}^{\alpha_{12},\beta_{12}}(e_{10};p_{12},q_{12};x_1,x_2) &= \frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1},\\
S_{n_{12},l}^{\alpha_{12},\beta_{12}}(e_{01};p_{12},q_{12};x_1,x_2) &= \frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2},\\
S_{n_{12},l}^{\alpha_{12},\beta_{12}}(e_{11};p_{12},q_{12};x_1,x_2) &= \frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2},\\
S_{n_{12},l}^{\alpha_{12},\beta_{12}}(e_{20};p_{12},q_{12};x_1,x_2) &= \frac{[n_1+l](p_1^{n_1+l-1}+2\alpha_1)x_1}{([n_1]+\beta_1)^2}+\frac{q_1[n_1+l][n_1+l-1]x_1^2}{([n_1]+\beta_1)^2}+\frac{\alpha_1^2}{([n_1]+\beta_1)^2},\\
S_{n_{12},l}^{\alpha_{12},\beta_{12}}(e_{02};p_{12},q_{12};x_1,x_2) &= \frac{[n_2+l](p_2^{n_2+l-1}+2\alpha_2)x_2}{([n_2]+\beta_2)^2}+\frac{q_2[n_2+l][n_2+l-1]x_2^2}{([n_2]+\beta_2)^2}+\frac{\alpha_2^2}{([n_2]+\beta_2)^2}.
\end{aligned}$$

Proof. From equation 2.1, we find that

$$\begin{aligned}
S_{n_{12},l}^{\alpha_{12}\beta_{12}}(e_{0,0};p_{12},q_{12};x_1,x_2) &= S_{n_1,l_1}^{\alpha_1,\beta_1}(e_0;p_1,q_1;x_1)S_{n_2,l_2}^{\alpha_2,\beta_2}(e_0;p_2,q_2;x_2),\\
S_{n_{12},l}^{\alpha_{12}\beta_{12}}(e_{1,0};p_{12},q_{12};x_1,x_2) &= S_{n_1,l_1}^{\alpha_1,\beta_1}(x_1;p_1,q_1;x_1)S_{n_2,l_2}^{\alpha_2,\beta_2}(e_0;p_2,q_2;x_2),\\
S_{n_{12},l}^{\alpha_{12}\beta_{12}}(e_{0,1};p_{12},q_{12};x_1,x_2) &= S_{n_1,l_1}^{\alpha_1,\beta_1}(e_0;p_1,q_1;x_1)S_{n_2,l_2}^{\alpha_2,\beta_2}(x_2;p_2,q_2;x_2),\\
S_{n_{12},l}^{\alpha_{12}\beta_{12}}(e_{1,1};p_{12},q_{12};x_1,x_2) &= S_{n_1,l_1}^{\alpha_1,\beta_1}(x_1;p_1,q_1;x_1)S_{n_2,l_2}^{\alpha_2,\beta_2}(x_2;p_2,q_2;x_2),\\
S_{n_{12},l}^{\alpha_{12}\beta_{12}}(e_{2,0};p_{12},q_{12};x_1,x_2) &= S_{n_1,l_1}^{\alpha_1,\beta_1}(x_1^2;p_1,q_1;x_1)S_{n_2,l_2}^{\alpha_2,\beta_2}(e_0;p_2,q_2;x_2),\\
S_{n_{12},l}^{\alpha_{12}\beta_{12}}(e_{0,2};p_{12},q_{12};x_1,x_2) &= S_{n_1,l_1}^{\alpha_1,\beta_1}(e_0;p_1,q_1;x_1)S_{n_2,l_2}^{\alpha_2,\beta_2}(x_2^2;p_2,q_2;x_2),
\end{aligned}$$

using these equalities, we can simply proof the Lemma 2.1. □

Lemma 2.2. *Let $S_{n_{12},l}^{\alpha_{12},\beta_{12}}(f;p_{12},q_{12},;x_1,x_2)$ be defined by 2.1. Then, we have*

$$\begin{aligned}
S_{n_{12},l}^{\alpha_{12}\beta_{12}}(((t_1-x_1)^2);p_{12},q_{12};x_1,x_2) &= \frac{[n_1+l][n_1+l-1]q_1-2[n_1+l]([n_1]+\beta_1)+([n_1]+\beta_1)^2}{([n_1]+\beta_1)^2}x_1^2\\
&\quad+\frac{[n_1+l](p_1^{n_1+l-1}+2\alpha_1)-2\alpha_1([n_1]+\beta_1)}{([n_1]+\beta_1)^2}x_1+\frac{\alpha_1^2}{(n+\beta_1)^2}.\\
S_{n_{12},l}^{\alpha_{12}\beta_{12}}(((t_2-x_2)^2);p_{12},q_{12};x_1,x_2) &= \frac{[n_2+l][n_2+l-1]q_2-2[n_2+l]([n_2]+\beta_2)+([n_2]+\beta_2)^2}{([n_2]+\beta_2)^2}x_2^2\\
&\quad+\frac{[n_2+l](p_2^{n_2+l-1}+2\alpha_2)-2\alpha_2([n_2]+\beta_2)}{([n_2]+\beta_2)^2}x_2+\frac{\alpha_2^2}{(n+\beta_2)^2}.
\end{aligned}$$

Proof. In view of Lemma 2.1 and linearity property, it is easy to prove Lemma 2.2. □

3. Main Results

Definition 3.1. Let $X,Y\subset\mathbb{R}$ be any two given intervals and the set $B(X\times Y)=\{f:X\times Y\to\mathbb{R}|\,f$ is bounded on $X\times Y\}$. For $f\in B(X\times Y)$, let the function $\omega_{total}(f;\cdot,*):[0,\infty)\times[0,\infty)\to\mathbb{R}$, defined for any $(\delta_1,\delta_2)\in[0,\infty)\times[0,\infty)$ by

$$\omega_{total}(f;\delta_1,\delta_2)=sup\{|f(x,y)-f(x',y')|:(x,y),(x',y')\in[0,\infty)\times[0,\infty),|x-x'|\le\delta_1,|y-y'|\le\delta_2\},$$

is called the first order modulus of smoothness of the function f or the total modulus of continuity of the function f.

In order to get the rate of convergence and degree of approximation for the operators $S_{n_{12},l}^{\alpha_{12}\beta_{12}}$, we consider $p_i=p_{n_i}$ and $q_i=q_{n_i}$ for $i=1,2$ such that $0<q_{n_i}<p_{n_i}\le 1$ satisfying

$$\lim_{n_i\to\infty}p_{n_i}^{n_i}\to a_i,\lim_{n_i\to\infty}q_{n_i}^{n_i}\to b_i \text{ and } \lim_{n_i\to\infty}p_{n_i}\to 1,\lim_{n_i\to\infty}q_{n_i}\to 1(i=1,2). \tag{3.1}$$

Here, we recall the following result due to Volkov [13]:

Theorem 3.1. *Let I and J be compact intervals of the real line. Let $L_{n_1,n_2}:C(I\times J)\to C(I\times J),(n_1,n_2)\in\mathbb{N}\times\mathbb{N}$ be linear positive operators. If*

$$\begin{aligned}
\lim_{n_1,n_2\to\infty}L_{n_1,n_2}(e_{ij}) &= e_{ij},(i,j)\in\{(0,0),(1,0),(0,1)\},\\
\lim_{n_1,n_2\to\infty}L_{n_1,n_2}(e_{20}+e_{02}) &= e_{20}+e_{02},
\end{aligned}$$

uniformly on $I\times J$, then the sequence $(L_{n_1,n_2}f)$ converges to f uniformly on $I\times J$ for any $f\in C(I\times J)$.

Theorem 3.2. *Let $e_{ij}(x_1,x_2)=x_1^i x_2^j (0\le i+j\le 2, i,j\in\mathbb{N})$ be the test functions defined on $I\times I$ and $(p_{n_i}),(q_{n_i}), i=1,2$ be the sequences defined by 3.1. If*

$$\lim_{n_1,n_2\to\infty}(S^{\alpha_{12}\beta_{12}}_{n_{12},l}e_{ij})(x_1,x_2)=e_{ij}(x_1,x_2),$$

uniformly on $I\times I$, then

$$\lim_{n_1,n_2\to\infty}(S^{\alpha_{12}\beta_{12}}_{n_{12},l}f)(x_1,x_2)=f(x_1,x_2).$$

uniformly for any $f\in C(I\times I)$.

Proof. Using the Theorem 3.1 and Lemma 2.1, Theorem 3.2 can easily be proved. □

Theorem 3.3. *[14] Let $L: C([0,\infty)\times[0,\infty))\to B([0,\infty)\times[0,\infty))$ be a linear positive operator. For any $f\in C(X\times Y)$, any $(x,y)\in X\times Y$ and any $\delta_1,\delta_2>0$, the following inequality*

$$\begin{aligned}|(Lf)(x,y)-f(x,y)| &\le |Le_{0,0}(x,y)-1||f(x,y)|+\Big[Le_{0,0}(x,y)+\delta_1^{-1}\sqrt{Le_{0,0}(x,y)(L(\cdot-x^2))^2(x,y)}\\ &+ \delta_2^{-1}\sqrt{Le_{0,0}(x,y)(L(*-y^2))^2(x,y)}+\delta_1^{-1}\delta_2^{-1}\sqrt{(Le_{0,0})^2(x,y)(L(\cdot-x^2))^2(x,y)(L(*-y^2))^2(x,y)}\Big]\\ &\times \omega_{total}(f;\delta_1,\delta_2),\end{aligned}$$

holds.

Theorem 3.4. *Let $f\in C(I\times I)$ and $(x_1,x_2)\in I\times I$. Then, for $(n_1,n_2)\in\mathbb{N}$ and for any $\delta_1,\delta_2>0$, we have*

$$|(S^{\alpha_{12}\beta_{12}}_{n_{12},l}f)(x_1,x_2)-f(x_1,x_2)|\le 4\omega_{total}(f;\delta_1,\delta_2),$$

where $\delta^{\alpha_{12}\beta_{12}}_{n_{12},l}(x_i)=\sqrt{S^{\alpha_{12}\beta_{12}}_{n_{12},l}(((t_i-x_i)^2);p_{12},q_{12};x_1,x_2)}$.

Proof Using Theorem 3.3 and Lemma 2.2 , we can arrive at the proof of the Theorem 3.4.

4. Local approximations

Let $C_B^2(I)=\{f\in C_B(I): f^{(i,j)}\in C_B(I), 1\le i,j\le 2\}$, where $C_B(I)$ is the space of all bounded and uniformly continuous functions on I and $f^{i,j}$ is $(i,j)^{th}$-order of partial derivative with respect to x,y of f, endowed with the norm

$$\|f\|_{C_B^2(I)}=\|f\|_{C_B(I)}+\sum_{i=1}^{2}\Big\|\frac{\partial^i f}{\partial x_1^i}\Big\|+\sum_{i=1}^{2}\Big\|\frac{\partial^i f}{\partial x_2^i}\Big\|.$$

The Peetre's K-functional of the function $f\in C_B(I)$ is given by

$$K(f;\delta)=\inf_{g\in C_B(I)^2}\{\|f-g\|_{C_B(I)}+\delta\|g\|_{C_B(I)^2},\delta>0\}. \tag{4.1}$$

The following inequality

$$K(f;\delta)\le M_1\{\omega_2(f;\sqrt{\delta})+min(1,\delta)\|f\|_{C_B(I)}\},$$

holds for all $\delta>0$ where M_1 is a constant independent of δ and f and $\omega_2(f;\sqrt{\delta})$ is the second order modulus of continuity which is defined in a similar manner as the second order modulus of continuity for one variable case

$$\omega(f;\sqrt{\delta})=\sup_{0<h\le\sqrt{\delta}}\sup_{x+2h\in[0,\infty)}|f(x+2h)-2f(x+h)+f(x)|.$$

Theorem 4.1. *Let (q_{n_i}) and (p_{n_i}) for $i=1,2$ are the real sequences defined in 3.1. Then, for $f\in C_B^2(I\times I)$, we have the following*

$$\begin{aligned}
|S_{n_{12},l}^{\alpha_{12}\beta_{12}}(g;x_1,x_2)-f(x_1,x_2)| &\leq 4K(f;M_{n_1,n_2}(x_1,x_2))\\
&+ \omega\left(f;\sqrt{\left(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}\right)^2+\left(\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\right)^2}\right)\\
&\leq M\left\{\omega_2\left(f:\sqrt{}(M_{n_1,n_2}(x_1,x_2))\right)+min\{1,M_{n_1,n_2}(x_1,x_2)\}\|f\|_{C_B^2(I)}\right\}\\
&+ \omega\left(f;\sqrt{\left(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}\right)^2+\left(\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\right)^2}\right),
\end{aligned}$$

where $M_{n_1,n_2}(x_1,x_2)=\left(\delta_{n_{12},l}^{\alpha_{12}\beta_{12}}(x_1)\right)^2+\left(\delta_{n_{12},l}^{\alpha_{12}\beta_{12}}(x_2)\right)^2$.

Proof. Consider the auxiliary operators

$$\widehat{S_{n_{12},l}^{\alpha_{12}\beta_{12}}}(f;x_1,x_2)=S_{n_{12},l}^{\alpha_{12}\beta_{12}}(f;x_1,x_2)-f\left(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1},\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\right)+f(x_1,x_2). \tag{4.2}$$

We get that,

$$\widehat{S_{n_{12},l}^{\alpha_{12}\beta_{12}}}(f;x_1,x_2)\leq 3\|f\|_{C_B(I)},\ \widehat{S_{n_{12},l}^{\alpha_{12}\beta_{12}}}(t_1-x_1;x_1,x_2)=0 \text{ and } \widehat{S_{n_{12},l}^{\alpha_{12}\beta_{12}}}(t_2-x_2;x_1,x_2)=0. \tag{4.3}$$

Let $g\in C_B^2(I)$ and $(x_1,x_2)\in I$. By the Taylor's theorem, we have

$$\begin{aligned}
g(u_1,u_2)-g(x_1,x_2) &= \frac{\partial g(x_1,x_2)}{\partial x_1}(u_1-x_1)+\int_{x_1}^{u_1}(u_1-\alpha_1)\frac{\partial^2 g(\alpha_1,x_1)}{\partial\alpha_1^2}d\alpha_1+\frac{\partial g(x_1,x_2)}{\partial x_2}(u_2-x_2)\\
&+\int_{x_2}^{u_2}(u_2-\alpha_2)\frac{\partial^2 g(\alpha_2,x_2)}{\partial\alpha_2^2}d\alpha_2.
\end{aligned} \tag{4.4}$$

Applying the auxiliary operator defined by 4.2 on both sides of 4.4, we find

$$\begin{aligned}
\widehat{S_{n_{12},l}^{\alpha_{12}\beta_{12}}}(g;x_1,x_2)-g(x_1,x_2) &= \widehat{S_{n_{12},l}^{\alpha_{12}\beta_{12}}}\left(\int_{x_1}^{u_1}(u_1-\alpha_1)\frac{\partial^2 g(\alpha_1,x_1)}{\partial\alpha_1^2}d\alpha_1;x_1,x_2\right)\\
&+\widehat{S_{n_{12},l}^{\alpha_{12}\beta_{12}}}\left(\int_{x_2}^{u_2}(u_2-\alpha_2)\frac{\partial^2 g(\alpha_2,x_2)}{\partial\alpha_2^2}d\alpha_2;x_1,x_2\right)\\
&= S_{n_{12},l}^{\alpha_{12}\beta_{12}}\left(\int_{x_2}^{u_2}(u_2-\alpha_2)\frac{\partial^2 g(\alpha_2,x_2)}{\partial\alpha_2^2}d\alpha_2;x_1,x_2\right)\\
&- \int_{x_1}^{\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}}\left(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}-\alpha_1\right)\frac{\partial^2 g(\alpha_1,x_1)}{\partial\alpha_1^2}d\alpha_1\\
&+ S_{n_{12},l}^{\alpha_{12}\beta_{12}}\left(\int_{x_2}^{u_2}(u_2-\alpha_2)\frac{\partial^2 g(\alpha_2,x_2)}{\partial\alpha_2^2}d\alpha_2;x_1,x_2\right)\\
&- \int_{x_2}^{\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}}\left(\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}-\alpha_2\right)\frac{\partial^2 g(\alpha_2,x_2)}{\partial\alpha_2^2}d\alpha_2.
\end{aligned}$$

Hence,

$$\begin{aligned}
|\widehat{S^{\alpha_{12}\beta_{12}}_{n_{12},l}}(g;x_1,x_2)-g(x_1,x_2)| &\leq S^{\alpha_{12}\beta_{12}}_{n_{12},l}\left(\Big|\int_{x_2}^{u_2}|u_2-\alpha_2|\Big|\frac{\partial^2 g(\alpha_2,x_2)}{\partial\alpha_2^2}\Big|d\alpha_2\Big|;x_1,x_2\right)\\
&+ \int_{x_1}^{\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}}\left|\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}-\alpha_1\right|\Big|\frac{\partial^2 g(\alpha_1,x_1)}{\partial\alpha_1^2}\Big|d\alpha_1\Big|\\
&+ S^{\alpha_{12}\beta_{12}}_{n_{12},l}\Big|\int_{x_2}^{u_2}|u_2-\alpha_2|\Big|\frac{\partial^2 g(\alpha_2,x_2)}{\partial\alpha_2^2}\Big|d\alpha_2\Big|;x_1,x_2\Big)\Big|\\
&+ \Big|\int_{x_2}^{\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}}\Big|\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}-\alpha_2\Big|\Big|\frac{\partial^2 g(\alpha_2,x_2)}{\partial\alpha_2^2}\Big|d\alpha_2\Big|\\
&\leq \Big\{S^{\alpha_{12}\beta_{12}}_{n_{12},l}((u_1-x_1)^2:x_1,x_2)+\Big(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}-x_1\Big)^2\Big\}\|g\|_{C_B^2(I)}\\
&+ \Big\{S^{\alpha_{12}\beta_{12}}_{n_{12},l}((u_2-x_2)^2:x_1,x_2)+\Big(\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}-x_2\Big)^2\Big\}\|g\|_{C_B^2(I)}\\
&= \{(\delta^{\alpha_{12}\beta_{12}}_{n_{12},l}(x_1))^2+(\delta^{\alpha_{12}\beta_{12}}_{n_{12},l}(x_2))^2\}\|g\|_{C_B^2(I)}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
|S^{\alpha_{12}\beta_{12}}_{n_{12},l}(g;x_1,x_2)-f(x_1,x_2)| &\leq |\widehat{S^{\alpha_{12}\beta_{12}}_{n_{12},l}}(f-g;x_1,x_2)|+|\widehat{S^{\alpha_{12}\beta_{12}}_{n_{12},l}}(g;x_1,x_2)-g(x_1,x_2)|+|g(x,y)-f(x,y)|\\
&+ \left|f\left(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1},\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\right)-f(x_1,x_2)\right|\\
&\leq 3\|f-g\|_{C_B(I)}+\|f-g\|_{C_B(I)}+|\widehat{S^{\alpha_{12}\beta_{12}}_{n_{12},l}}(g;x_1,x_2)-g(x_1,x_2)|\\
&+ \left|f\left(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1},\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\right)-f(x_1,x_2)\right|\\
&\leq 4\|f-g\|_{C_B(I)}+\{(\delta^{\alpha_{12}\beta_{12}}_{n_{12},l}(x_1))^2+(\delta^{\alpha_{12}\beta_{12}}_{n_{12},l}(x_2))^2\}\|g\|_{C_B^2(I)}\\
&+ \left|f\left(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1},\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\right)-f(x_1,x_2)\right|\\
&\leq 4\|f-g\|_{C_B(I)}+2M_{n_1,n_2}(x_1,x_2)\|_{C_B^2(I)}+\omega\left(f;\sqrt{\Big(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}\Big)^2+\Big(\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\Big)^2}\right).
\end{aligned}$$

Next, using the equation 4.1, we get

$$\begin{aligned}
|S^{\alpha_{12}\beta_{12}}_{n_{12},l}(g;x_1,x_2)-f(x_1,x_2)| &\leq 4K(f;M_{n_1,n_2}(x_1,x_2))\\
&+ \omega\left(f;\sqrt{\Big(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}\Big)^2+\Big(\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\Big)^2}\right)\\
&\leq M\left\{\omega_2\Big(f:\sqrt{(}M_{n_1,n_2}(x_1,x_2))\Big)+min\{1,M_{n_1,n_2}(x_1,x_2)\}\|f\|_{C_B^2(I)}\right\}\\
&+ \omega\left(f;\sqrt{\Big(\frac{[n_1+l]x_1+\alpha_1}{[n_1]+\beta_1}\Big)^2+\Big(\frac{[n_2+l]x_2+\alpha_2}{[n_2]+\beta_2}\Big)^2}\right).
\end{aligned}$$

□

Now, we discuss the degree of approximation for the (p,q)-Bivariate-Schurer-Stancu operators in the Lipschitz class. We define the Lipschitz class $Lip_M(\gamma_1,\gamma_2)$ by means of two variables as follows:

$$|f(t_1,t_2)-f(x_1,x_2)|\leq M|t_1-x_1|^{\gamma_1}|t_2-x_2|^{\gamma_2},$$

where $0 < \gamma_1, \gamma_2 \leq 1$ and for any $(t_1,t_2),(x_1,x_2) \in I \times I$.

Theorem 4.2. *Let $f \in Lip_M(\gamma_1,\gamma_2)$ and $(q_{n_i}),(p_{n_i})$, $i = 1,2$ are defined in 3.1. Then for all $(x_1,x_2) \in I \times I$, we have*

$$|S_{n_{12},l}^{\alpha_{12}\beta_{12}}(g;x_1,x_2) - f(x_1,x_2)| \leq M\delta_{n_1}^{\gamma_1/2}(x_1)\delta_{n_2}^{\gamma_2/2}(x_2)$$

where $\delta_{n_i}(x_i) = S_{n_{12},l}^{\alpha_{12}\beta_{12}}(((t_i - x_i)^2);p_{12},q_{12};x_1,x_2)$.

Proof. Since $f \in Lip_M(\gamma_1,\gamma_2)$, we can write

$$\begin{aligned}
|S_{n_{12},l}^{\alpha_{12}\beta_{12}}(f;q_{n_{12}},p_{n_{12}};x_1,x_2) - f(x_1,x_2)| &\leq S_{n_{12},l}^{\alpha_{12}\beta_{12}}(|f(t_1,t_2) - f(x_1,x_2)|;q_{n_{12}},p_{n_{12}};x_1,x_2) \\
&\leq MS_{n_{12},l}^{\alpha_{12}\beta_{12}}(|t_1 - x_1|^{\gamma_1}|t_2 - x_2|^{\gamma_2};q_{n_{12}},p_{n_{12}};x_1,x_2) \\
&= MS_{n_{12},l}^{\alpha_{12}\beta_{12}}(|t_1 - x_1|^{\gamma_1};q_{n_{12}},p_{n_{12}};x_1,x_2)S_{n_{12},l}^{\alpha_{12}\beta_{12}}(|t_2 - x_2|^{\gamma_2};q_{n_{12}},p_{n_{12}};x_1,x_2).
\end{aligned}$$

Next, we use the Holder inequality with $p = \frac{2}{\gamma_1}$, $q = \frac{2}{2-\gamma_1}$ and $p = \frac{2}{\gamma_1}$, $q = \frac{2}{2-\gamma_2}$, respectively, we have

$$\begin{aligned}
|S_{n_{12},l}^{\alpha_{12}\beta_{12}}(f;q_{n_{12}},p_{n_{12}};x_1,x_2) - f(x_1,x_2)| &\leq \left\{S_{n_{12},l}^{\alpha_{12}\beta_{12}}((t_1 - x_1)^2;q_{n_{12}},p_{n_{12}};x_1,x_2)\right\}^{\frac{\gamma_1}{2}}\left\{S_{n_{12},l}^{\alpha_{12}\beta_{12}}((1;q_{n_{12}},p_{n_{12}};x_1,x_2)\right\}^{\frac{2}{2-\gamma_1}} \\
&\times \left\{S_{n_{12},l}^{\alpha_{12}\beta_{12}}((t_2 - x_2)^2;q_{n_{12}},p_{n_{12}};x_1,x_2)\right\}^{\frac{\gamma_2}{2}}\left\{S_{n_{12},l}^{\alpha_{12}\beta_{12}}((1;q_{n_{12}},p_{n_{12}};x_1,x_2)\right\}^{\frac{2}{2-\gamma_2}} \\
&= M\delta_{n_1}^{\gamma_1/2}(x_1)\delta_{n_2}^{\gamma_2/2}(x_2).
\end{aligned}$$

□

References

References

[1] F. Schurer, Positive linear operators in approximation theory, Mathematical Institute of the technological University Delft, Report (1962).
[2] A. Lupas, A q-analogue of the Bernstein operators, Seminar on Numerical and Statistical Calculus, University of Cluj-Napoca, 9 (1987) 85-98.
[3] G. M. Philips, Bernstein polynomials based on the q-integers, The heritage of P.L. Chebyshev, Ann. Numer. Math., 4 (1997) 511-518.
[4] A. Aral, V. Gupta, R.P. Agarwal, Applications of q-Calculus in Operator Theory, Springer Science+Business Media New York, 2013.
[5] M. Mursaleen, Khursheed J. Ansari, Asif Khan, On (p,q)-analogue of Bernstein operators, Appl. Math. Comput., 266 (2015) 874-882.
[6] R. Chakarabarti and R. Jagannathan, A (p,q)-oscillator realization of two parameter quantum algebras, J. Phys. A: Math. Gen., 24 (1991) 711-718.
[7] M. Mursaleen, Khursheed J. Ansari, Asif Khan, Some approximation results by (p,q)-analogue of Bernstein -Stancu operators, Appl. Math. Comput., 264 (2015) 392-402.
[8] Mursaleen, Mohammad; Nasiruzzaman, Md; Nurgali, Ashirbayev Some approximation results on Bernstein-Schurer operators defined by (p,q)-integers. J. Inequal. Appl. 2015, 2015:2
[9] A. Wafi, N. Rao; Approximation properties of (p;q)-variant of Stancu-Schurer operators, arXiv:1508.01852v3 [math.NT] 31 Dec 2015.
[10] V. N. Mishra, M. Mursaleen, S. Pandey; Approximation properties of Chlodowsky variant of (p,q) Bernstein-Stancu-Schurer operators, arXiv:1508.06888v3 [math.CA] 26 Nov 2015.
[11] A. Tucer; (p,q)-Generalization of SzszMirakyan operators, Math. Meth. Appl. Sci., DOI: 10.1002/mma.3721.
[12] V. Gupta; (p,q)-Szsz-Mirakjan-Baskakov operators, Complex Analysis and Operator Theory, 12/2015; DOI:10.1007/s11785-015-0521-4
[13] V. I. Volkov, On the convergence of sequences of linear positive operators in the space of continuous functions of two variables (in Russian), Dokl. Akad. Nauk SSSR (NS) 115 (1957) 17-19.
[14] F. Stancu, Approximarca funcțiilor de două şi mai multe vsrabile prin șiruri de operatori liniari și positivi, PhD Thesis, Cluj-Napoca, 1984 (Romanian).